

OK TO ENTER: /C.Y.W./

09/09/2009

IN THE SPECIFICATION:

Please amend paragraph [0009] as follows:

[0009] In one embodiment, an isolated polypeptide comprising the amino acid sequence Y (Trp/Phe) Xaa₁ Xaa₂ Xaa₃ Xaa₄ Xaa₅ (Trp/Phe) Xaa₆ Xaa₇ (Trp/Phe) Z (SEQ ID NOs:17-20) is provided. Y, which may or may not be present, is a peptidic structure containing at least one cysteine residue and having the formula (Xaa)_n. Xaa is any amino acid residue and n is an integer from 1 to 20. Z, which may or may not be present, is a peptidic structure containing at least one cysteine residue and having the formula (Xaa)_n, wherein Xaa is any amino acid residue and n is an integer from 1 to 20. The amino acid residues of in Xaa₁ through Xaa₇ can be any amino acid and the amino acid residues of Xaa₁ through Xaa₅ are positively charged.

Please amend paragraph [0010] as follows:

[0010] In another embodiment, an isolated polypeptide comprising the amino acid sequence Y (Trp/Phe) Xaa₁ Xaa₂ Xaa₃ Xaa₄ Xaa₅ (Trp/Phe) Xaa₆ Xaa₇ Xaa₈ (Trp/Phe) Z (SEQ ID NOs:21-24) is provided. Y, which may or may not be present, is a peptidic structure containing at least one cysteine residue and having the formula (Xaa)_n. Xaa is any amino acid residue and n is an integer from 1 to 20. Z, which may or may not be present, is a peptidic structure containing at least one cysteine residue and having the formula (Xaa)_n, wherein Xaa is any amino acid residue and n is an integer from 1 to 20. The amino acid residues of Xaa₁ through Xaa₈ is any amino acid, and at least two of the residues of Xaa₁ through Xaa₅ are positively charged.